



Autolus Therapeutics' CAR T Therapy AUCATZYL® (Obecabtagene Autoleucel) Granted European Marketing Authorization for Adult Patients (age 26 and older) with Relapsed or Refractory B-Cell Precursor Acute Lymphoblastic Leukemia (R/R B-ALL)

July 21, 2025 at 7:00 AM EDT

- Approval is based on FELIX clinical trial of obecabtagene autoleucel (obe-cel) in adult patients with r/r B-ALL, demonstrating high and durable response rates and low toxicity
- EC approval follows positive CHMP opinion, MHRA conditional marketing authorization and FDA approval

LONDON and GAITHERSBURG, Md., July 21, 2025 (GLOBE NEWSWIRE) -- Autolus Therapeutics plc (Nasdaq: AUTL), an early commercial-stage biopharmaceutical company developing, manufacturing and delivering next-generation programmed T cell therapies and candidates, announces today that the European Commission (EC) has granted marketing authorization for AUCATZYL® (obecabtagene autoleucel or "obe-cel") for the treatment of adult patients, 26+, with relapsed or refractory B-cell precursor acute lymphoblastic leukemia (r/r B-ALL).

The EC approval was based on the results of the FELIX study, an open-label, multi centre, single arm study in adult patients with relapsed or refractory B-cell acute lymphoblastic leukaemia. The results were published in the New England Journal of Medicine in November 2024¹. In the pivotal cohort of patients, (cohort IIA (n=94)), the Complete Response/Complete Response with Incomplete Haematological Recovery (CR/CRi) for patients who received at least one infusion of obecabtagene autoleucel was 76.6%. Median response duration for all infused patients was 21.2 months. Median event-free survival (EFS) was 11.9 months and the estimated 6- and 12-month event-free survival rates were 65.4% and 49.5%, respectively.

The most common non-laboratory Grade 3 or higher adverse reactions were infections-pathogen unspecified (32%), febrile neutropenia (24%) and bacterial infectious disorders (11%). Cytokine release syndrome developed in 87 of the 127 patients (68.5%), with events of grade 3 or higher in three patients (2.4%). Immune effector cell-associated neurotoxicity syndrome developed in 29 of the 127 patients (22.8%), with grade 3 or higher occurring in nine patients (7%).

"We believe AUCATZYL represents an important new treatment option for physicians treating adult r/r B-ALL patients. With the EU marketing authorization, we are now evaluating market entry opportunities in EU countries," said **Dr. Christian Itin, Chief Executive Officer of Autolus**.

Obe-cel is an autologous CD19 CAR T cell therapy with a proprietary CD19 CAR, invented by a team led by Dr. Martin Pule, at University College London, along with collaborators at Great Ormond Street Hospital and University College London Hospital. The CAR is designed to have a fast "off-rate" which mimics physiological T-cell receptor interactions².

ALL is an aggressive type of blood cancer that can also involve the lymph nodes, spleen, liver, central nervous system and other organs. In Europe, there are approximately 6,000² new cases of ALL diagnosed every year. In frontline treatment for adult B-ALL, up to 50% of patients will ultimately relapse³. Survival rates remain very poor in adult patients with r/r ALL, with median overall survival of eight months with conventional treatments⁴, and the standard-of-care treatment can trigger severe toxicities⁵.

The EC approval applies to all 27 European Union Member States, Iceland, Norway and Liechtenstein. AUCATZYL is currently approved by the U.S. Food and Drug Administration (FDA) and authorized by the U.K. Medicines and Healthcare products Regulatory Agency (MHRA).

References

1. Roddie C, et al "[Obcabtagene autoleucel in B-cell acute lymphoblastic leukemia](#)" *N Engl J Med* 2024; DOI: 10.1056/NEJMoa2406526
2. <https://www.sciencedirect.com/science/article/pii/S0006497120310946?via%3Dihub>
3. Cancer Research UK - <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/leukaemia-all/incidence>
4. Aureli A, Marziani B, Venditti A, Sconocchia T, Sconocchia G. Acute lymphoblastic leukemia immunotherapy treatment: now, next, and beyond. *Cancers (Basel)*. 2023;15:3346.
5. Dhakal P, Kaur J, Gundabolu K, Bhatt VR. Immunotherapeutic options for management of relapsed or refractory B-cell acute lymphoblastic leukemia: how to select newly approved agents? *Leuk Lymphoma*. 2020;61:7-17.

About Autolus Therapeutics plc

Autolus Therapeutics plc (Nasdaq: AUTL) is an early commercial-stage biopharmaceutical company developing, manufacturing and delivering next-generation T cell therapies and candidates for the treatment of cancer and autoimmune disease. Using a broad suite of proprietary and modular T cell programming technologies, Autolus is engineering precisely targeted and controlled T cell therapies that are designed to better recognize target cells, break down their defense mechanisms and eliminate these cells. Autolus has a marketed therapy, AUCATZYL®, and a pipeline of product candidates in development for the treatment of hematological malignancies, solid tumors and autoimmune diseases. For more information, please visit www.autolus.com.

About obe-cel FELIX clinical trial

Autolus' Phase 1b/2 clinical trial of obe-cel enrolled adult patients with r/r B-precursor ALL. The trial had a Phase 1b component prior to proceeding to the single arm, Phase 2 clinical trial. The primary endpoint in the pivotal cohort was overall response rate, and the secondary endpoints included duration of response, MRD negative complete remission rate and safety. The trial enrolled over 100 patients across 30 of the leading academic and non-academic centers in the United States, United Kingdom and Europe. [NCT04404660].

U.S. Indication and Safety Information

INDICATION

AUCATZYL[®] is a CD19-directed genetically modified autologous T cell immunotherapy indicated for the treatment of adult patients with relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL).

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME, NEUROLOGIC TOXICITIES, and SECONDARY HEMATOLOGICAL MALIGNANCIES

- **Cytokine Release Syndrome (CRS) occurred in patients receiving AUCATZYL. Do not administer AUCATZYL to patients with active infection or inflammatory disorders. Prior to administering AUCATZYL, ensure that healthcare providers have immediate access to medications and resuscitative equipment to manage CRS.**
- **Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS), including fatal and life-threatening reactions, occurred in patients receiving AUCATZYL, including concurrently with CRS or after CRS resolution. Monitor for neurologic signs and symptoms after treatment with AUCATZYL. Prior to administering AUCATZYL, ensure that healthcare providers have immediate access to medications and resuscitative equipment to manage neurologic toxicities. Provide supportive care and/or corticosteroids, as needed.**
- **T cell malignancies have occurred following treatment of hematologic malignancies with BCMA- and CD19-directed genetically modified autologous T cell immunotherapies.**

WARNINGS AND PRECAUTIONS

Cytokine Release Syndrome (CRS)

Cytokine Release Syndrome (CRS) occurred following treatment with AUCATZYL. CRS was reported in 75% (75/100) of patients including Grade 3 CRS in 3% of patients. The median time to onset of CRS was 8 days following the first infusion (range: 1 to 23 days) with a median duration of 5 days (range: 1 to 21 days). The most common manifestations of CRS included fever (100%), hypotension (35%), and hypoxia (19%).

Prior to administering AUCATZYL, ensure that healthcare providers have immediate access to medications and resuscitative equipment to manage CRS. During and following treatment with AUCATZYL, closely monitor patients for signs and symptoms of CRS daily for at least 14 days at the healthcare facility following the first infusion. Continue to monitor patients for CRS for at least 4 weeks following each infusion with AUCATZYL. Counsel patients to seek immediate medical attention should signs or symptoms of CRS occur at any time. At the first sign of CRS, immediately evaluate the patient for hospitalization and institute treatment with supportive care based on severity and consider further management per current practice guidelines.

Neurologic Toxicities

Neurologic toxicities including Immune Effector Cell-associated Neurotoxicity Syndrome (ICANS), which were fatal or life-threatening, occurred following treatment with AUCATZYL. Neurologic toxicities were reported in 64% (64/100) of patients, including Grade ≥ 3 in 12% of patients. The median time to onset of neurologic toxicities was 10 days (range: 1 to 246 days) with a median duration of 13 days (range: 1 to 904 days). Among patients with neurologic toxicities, the most common symptoms ($> 5\%$) included ICANS (38%), headache (34%), encephalopathy (33%), dizziness (22%), tremor (13%), anxiety (9%), insomnia (9%), and delirium (8%).

Immune Effector Cell-associated Neurotoxicity Syndrome (ICANS)

ICANS events occurred in 24% (24/100) of patients, including Grade ≥ 3 in 7% (7/100) of patients. Of the 24 patients who experienced ICANS, 33% (8/24) experienced an onset after the first infusion, but prior to the second infusion of AUCATZYL.

The median time to onset for ICANS events after the first infusion was 8 days (range: 1 to 10 days) and 6.5 days (range: 2 to 22 days) after the second infusion, with a median duration of 8.5 days (range: 1 to 53 days).

Eighty-eight percent (21/24) of patients received treatment for ICANS. All treated patients received high-dose corticosteroids and 42% (10/24) of patients received anti-epileptics prophylactically. Prior to administering AUCATZYL, ensure that healthcare providers have immediate access to medications and resuscitative equipment to manage ICANS.

Counsel patients to seek medical attention should signs or symptoms of neurologic toxicity/ ICANS occur. At the first sign of Neurologic Toxicity /ICANS, immediately evaluate patients for hospitalization and institute treatment with supportive care based on severity and consider further management per current practice guidelines.

Effect on Ability to Drive and Use Machines

Due to the potential for neurologic events, including altered mental status or seizures, patients receiving AUCATZYL are at risk for altered or decreased consciousness or coordination in the eight weeks following AUCATZYL infusion or until resolution of the neurological event by the treating physician. Advise patients to refrain from driving and engaging in hazardous occupations or activities, such as operating heavy or potentially dangerous machinery, during this initial period.

Prolonged Cytopenias

Patients may exhibit cytopenias including anemia, neutropenia, and thrombocytopenia for several weeks after treatment with lymphodepleting chemotherapy and AUCATZYL. In patients who were responders to AUCATZYL, Grade ≥ 3 cytopenias that persisted beyond Day 30 following AUCATZYL infusion were observed in 71% (29/41) of patients and included neutropenia (66%, 27/41) and thrombocytopenia (54%, 22/41). Grade 3 or

higher cytopenias that persisted beyond Day 60 following AUCATZYL infusion was observed in 27% (11/41) of patients and included neutropenia (17%, 7/41) and thrombocytopenia (15%, 6/41). Monitor blood counts after AUCATZYL infusion.

Infections

Severe, including life-threatening and fatal infections occurred in patients after AUCATZYL infusion. Non-COVID-19 infections of all grades occurred in 67% (67/100) of patients. Grade 3 or higher non-COVID-19 infections occurred in 41% (41/100) of patients. AUCATZYL should not be administered to patients with clinically significant active systemic infections. Monitor patients for signs and symptoms of infection before and after AUCATZYL infusion and treat appropriately. Administer prophylactic antimicrobials according to local guidelines.

Grade 3 or higher febrile neutropenia was observed in 26% (26/100) of patients after AUCATZYL infusion and may be concurrent with CRS. In the event of febrile neutropenia, evaluate for infection and manage with broad-spectrum antibiotics, fluids, and other supportive care as medically indicated.

Viral reactivation, potentially severe or life-threatening, can occur in patients treated with drugs directed against B cells. There is no experience with manufacturing AUCATZYL for patients with a positive test for human immunodeficiency virus (HIV) or with active hepatitis B virus (HBV) or active hepatitis C virus (HCV). Perform screening for HBV, HCV and HIV in accordance with clinical guidelines before collection of cells for manufacturing.

Hypogammaglobulinemia

Hypogammaglobulinemia and B-cell aplasia can occur in patients after AUCATZYL infusion. Hypogammaglobulinemia was reported in 10% (10/100) of patients treated with AUCATZYL including Grade 3 events in 2 patients (2%).

Immunoglobulin levels should be monitored after treatment with AUCATZYL and managed per institutional guidelines including infection precautions, antibiotic or antiviral prophylaxis, and immunoglobulin replacement.

The safety of immunization with live viral vaccines during or following treatment with AUCATZYL has not been studied. Vaccination with live viral vaccines is not recommended for at least 6 weeks prior to the start of lymphodepleting chemotherapy treatment, during AUCATZYL treatment, and until immune recovery following treatment with AUCATZYL.

Hemophagocytic Lymphohistiocytosis/Macrophage Activation Syndrome (HLH/MAS)

HLH/MAS including fatal and life-threatening reactions occurred after treatment with AUCATZYL. HLH/MAS was reported in 2% (2/100) of patients and included Grade 3 and Grade 4 events with a time of onset at Day 22 and Day 41, respectively. One patient experienced a concurrent ICANS events after AUCATZYL infusion and died due to sepsis with ongoing HLH/MAS that had not resolved. Administer treatment for HLH/MAS according to institutional standards.

Hypersensitivity Reactions

Serious hypersensitivity reactions, including anaphylaxis, may occur due to dimethyl sulfoxide (DMSO), an excipient used in AUCATZYL. Observe patients for hypersensitivity reactions during and after AUCATZYL infusion.

Secondary Malignancies

Patients treated with AUCATZYL may develop secondary malignancies. T cell malignancies have occurred following treatment of hematologic malignancies with BCMA- and CD19-directed genetically modified autologous T cell immunotherapies. Mature T cell malignancies, including CAR-positive tumors, may present as soon as weeks following infusion, and may include fatal outcomes. Monitor lifelong for secondary malignancies. In the event that a secondary malignancy occurs, contact Autolus at 1-855-288-5227 for reporting and to obtain instructions on the collection of patient samples for testing.

Adverse Reactions

The safety of AUCATZYL was evaluated in the FELIX study in which 100 patients with relapsed or refractory B-cell acute lymphoblastic leukemia (B-ALL) received AUCATZYL at a median dose of 410×10^6 CD19 CAR-positive viable T cells (range: 10 to 480×10^6 CD19 CAR-positive viable T cells with 90% of patients receiving the recommended dose of $410 \times 10^6 \pm 25\%$).

The most common serious adverse reactions of any Grade (incidence $\geq 2\%$) included infections-pathogen unspecified, febrile neutropenia, ICANS, CRS, fever, bacterial infectious disorders, encephalopathy, fungal infections, hemorrhage, respiratory failure, hypotension, ascites, HLH/MAS, thrombosis and hypoxia. Nine patients (9%) experienced fatal adverse reactions which included infections (sepsis, pneumonia, peritonitis), ascites, pulmonary embolism, acute respiratory distress syndrome, HLH/MAS and ICANS. Of the 9 patients, five patients who died from infections had pre-existing and ongoing neutropenia prior to receiving bridging therapy, lymphodepletion chemotherapy treatment and/or AUCATZYL.

Please see full [Prescribing Information](#), including **BOXED WARNING** and Medication Guide.

Forward Looking Statements

This press release contains forward-looking statements within the meaning of the "safe harbor" provisions of the Private Securities Litigation Reform Act of 1995. Forward-looking statements are statements that are not historical facts, and in some cases can be identified by terms such as "may," "will," "could," "expects," "plans," "anticipates," and "believes." These risks and uncertainties include, but are not limited to the impact of worsening macroeconomic conditions on Autolus' business, financial position, strategy and anticipated milestones, including Autolus' ability to conduct ongoing and planned clinical trials; Autolus' ability to obtain a clinical supply of current or future product candidates or commercial supply of obe-cel or any future approved products; Autolus' ability to obtain and maintain regulatory approval of its product candidates, including obe-cel and potential expansions into additional indications; Autolus' ability and plans in continuing to establish and expand a commercial infrastructure and to successfully launch, market and sell obe-cel and any future approved products; Autolus' ability to obtain marketing approval for obe-cel in additional geographies in the future; the delay of any current or planned clinical trials, whether due to patient enrollment delays or otherwise; Autolus' ability to successfully demonstrate the safety and efficacy of its product candidates and gain approval of its product candidates, including obe-cel and potential expansions into additional indications; Autolus' ability and plans in continuing to establish and expand a commercial infrastructure and to successfully launch, market and sell obe-cel and any future approved products; Autolus' ability to obtain marketing approval for obe-cel in additional geographies in the future; the delay of any current or planned clinical trials, whether due to patient enrollment delays or otherwise; Autolus' ability to successfully demonstrate the safety and efficacy of its product candidates and gain approval of its product candidates on a timely basis, if at all; competition with respect to market opportunities; and possible safety and efficacy concerns. For a discussion of other risks and uncertainties, and other important factors, any of which could cause Autolus' actual results to differ from those contained in the forward-looking statements, see the section titled "Risk Factors" in Autolus' Annual Report on Form 10-K filed with the Securities and Exchange Commission, or the SEC, on March 20, 2025 as well as discussions of potential risks, uncertainties, and other important factors in Autolus' subsequent filings with the Securities and Exchange Commission.

All information in this press release is as of the date of the release, and Autolus undertakes no obligation to publicly update any forward-looking statement, whether as a result of new information, future events, or otherwise, except as required by law. You should, therefore, not rely on these forward-looking statements as representing Autolus' views as of any date subsequent to the date of this press release.

Contact:

Amanda Cray
+1 617-967-0207
a.cray@autolus.com