


Autolus

Nasdaq: AUTL



December 2019

Autolus AACR Data Update
June 2020

Disclaimer

These slides and the accompanying oral presentation contain forward-looking statements within the meaning of the “safe harbor” provisions of The Private Securities Litigation Reform Act of 1995, including statements about the Company's plans to develop and commercialize its product candidates, the Company's ongoing and planned clinical trials, including the timing and initiation of such trials and statements regarding whether or not such trials will be considered pivotal trials, the anticipated benefits of the Company's financial condition and results of operations, including its expected cash runway; the development of Autolus' product candidates, including statements regarding the timing of initiation, completion and the outcome of pre-clinical studies or clinical trials and related preparatory work, and the periods during which the results of the studies and trials will become available; Autolus' plans to research, develop, manufacture and commercialize its product candidates; the potential for Autolus' product candidates to be alternatives in the therapeutic areas investigated; and Autolus' manufacturing capabilities and strategy. All statements other than statements of historical fact contained in this presentation, including statements regarding the Company's future results of operations and financial position, business strategy and plans and objectives of management for future operations, are forward-looking statements. In some cases, you can identify forward-looking statements by terms such as “may,” “should,” “expects,” “plans,” “anticipates,” “could,” “intends,” “target,” “projects,” “contemplates,” “believes,” “estimates,” “predicts,” “potential” or “continue” or the negative of these terms or other similar expressions. These statements involve known and unknown risks, uncertainties and other important factors that may cause the Company's actual results, performance or achievements to be materially different from any future results, performance or achievements expressed or implied by the forward-looking statements. Factors that may cause actual results to differ materially from any future results expressed or implied by any forward looking statements include the risks described in the “Risk Factors” section of the Company's Annual Report on Form 20-F for the year ended December 31, 2019, as well as those set forth from time to time in the Company's other SEC filings, available at www.sec.gov. The forward-looking statements contained in this presentation reflect the Company's views as of the date of this presentation regarding future events, except as required by law, and the Company does not assume any obligation to update any forward-looking statements. You should, therefore, not rely on these forward-looking statements as representing the Company's views as of any date subsequent to the date of this presentation.

Certain data in this presentation was obtained from various external sources. Such data speak only as of the date referenced in this presentation and neither the Company nor its affiliates, advisors or representatives make any representation as to the accuracy or completeness of that data or undertake to update such data after the date of this presentation. Such data involve risks and uncertainties and are subject to change based on various factors.

Agenda

1. Welcome and Introduction: Dr. Christian Itin, Chairman and CEO
2. T Cell Lymphoma Program:
 - AUTO5 Data Review: Dr. Mathieu Ferrari, Associate Director of Binder Discovery
3. Autolus Solid Tumor Programs:
 - AUTO6NG Data Review: Dr. Muhammad Al-Hajj, SVP, Head of Translational Medicine
 - AUTO7 Data Review: Dr. Marco Della Peruta, Senior Scientist, Immunobiology
4. Summary & Next Steps: Dr. Christian Itin
5. Q&A: Dr. Christian Itin, Dr. Martin Pule (Founder & SVP, CSO), Dr. Vijay Reddy (SVP, CMO), Dr. Muhammad Al-Hajj, Dr. Mathieu Ferrari and Dr. Marco Della Peruta

Welcome and introduction

Dr. Christian Itin

Chairman and CEO

Broad expertise in translational research and binder discovery



Dr. Christian Itin

Chairman & CEO

Previously CEO of Micromet; led development of Blincyto[®], the first FDA-approved redirected T cell therapy



Dr. Muhammad Al-Hajj

SVP, Head of Translational Medicine

Experienced drug hunter & cancer biologist. 15+ years Oncology R&D leadership, GlaxoSmithKline, Novartis, AstraZeneca



Dr. Martin Pule

Founder & SVP, CSO

Founder of Autolus; World leading expert in the development of CAR T cell therapies; Clinical senior lecturer & hon. consultant at UCL; Fulbright at Baylor



Dr. Marco Della Peruta

Senior Scientist, Immunobiology

Human Oncological Pathology at University of Verona, School of Medicine, Italy. Previously, lecturer in Haematology at UCL Cancer Institute, UCL, London, UK



Dr. Vijay Peddareddigari

SVP, CMO

Experienced oncologist and drug developer; MD Anderson, GSK and most recently J&J



Dr. Mathieu Ferrari

Associate Director of Binder Discovery

Experimental Medicine, Queen Mary University of London, UK. Previously, post-doctoral research fellow at Centre for Experimental Medicine and Rheumatology in Queen Mary University of London

Broad pipeline of next generation programs


Designed to address limitations of current T cell therapies

Product	Indication	Target	Pre-clinical	Phase 1
B Cell Malignancies				
AUTO1NG	ALL	CD19 & CD22		H2 2020
AUTO3NG	DLBCL	CD19 & CD22		Life cycle mgmt
T Cell Lymphoma				
AUTO5	TRBC2+ Peripheral TCL	TRBC2		H1 2021
GD2+ Tumors				
AUTO6NG	Neuroblastoma; Melanoma; Osteosarcoma; SCLC	GD2		H1 2021
Prostate Cancer				
AUTO7	Prostate Cancer	PSMA		H1 2021
Multiple Myeloma				
AUTO8	Multiple Myeloma	BCMA & CAR X		H2 2020

NG = Next Generation, SCLC = Small Cell Lung Cancer

A broad toolkit building on our core strategy of modular innovation

Advanced T cell programming





T Cell Lymphoma

AUTO5 – tailored for T Cell Lymphoma

T Cell Lymphoma


No standard of care after first relapse

There is currently no T cell therapy approved for T cell lymphoma

- T cell lymphoma is an aggressive disease with a very poor prognosis for patients
- Median 5 yrs OS: 32%
- Standard of care is variable and often based on high-dose chemotherapy and stem cell transplants
- A large portion of T cell lymphoma patients are refractory to or relapse following treatment with standard therapies
- T cell lymphomas have not benefited from advances in immunotherapeutic approaches




Addressing T cell lymphomas

Three key elements - AUTO4, AUTO5 and a companion diagnostic test





Challenges Targeting TCR β

Differences between TRBC1 and TRBC2 are small




		NK-KN 4/5		F-Y 36
TRBC1	1	EDLNKVFPPPEVAVFEPSEAEISHTQKATLVCLATGFF		PDHVELSWVWNGK
TRBC2	1	EDLNKVFPPPEVAVFEPSEAEISHTQKATLVCLATGFF		PDHVELSWVWNGK
TRBC1	51	EVHSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRQCQVQF		
TRBC2	51	EVHSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRQCQVQF		
TRBC1	101	YGLSENDEWTODRAKPVTQIVSAEAWGRADCGFTSV		SYOOGVLSAT
TRBC2	101	YGLSENDEWTODRAKPVTQIVSAEAWGRADCGFTSE		SYOOGVLSAT
			V-E 135	








Addressing T cell lymphomas

Three key elements - AUTO4, AUTO5 and a companion diagnostic test



Analysis of α TRBC1-TCR contact interface (Crystallized complex 2.4Å)


α TRBC1 binds the TCR via specific contact with TCR β 1 chain in close proximity to CD3 ϵ



Interactions are driven via HCDR1 and HCDR3 with residues Asn119 and Lys120 on TCR β 1

Structure guided in-silico mutagenesis



Molecular modelling identifies 3 key point mutations enhancing TRBC2 specificity



Mut3 (α TRBC2) shows an inversion of TCR specificity

CAR T cell *in-vitro* characterization



Optimized spacer/endodomain demonstrates specific cytotoxicity towards TRBC2 cells



α TRBC2 CAR efficiently kills TRBC2⁺ PBMCs at endogenous levels of TCR

CAR T cell *in-vitro* characterization

α TRBC2 CAR T cells show enhanced cytokine production and proliferative capacity




2-way Anova

Significant proliferation and cytokine release for α TRBC2 CAR against TRBC2⁺ target cells

TRBC2 in-vivo CAR Activity

Selectivity of CAR targeting confirmed in mixed tumor NSG model



α TRBC2 CAR efficiently clears TRBC2+ targets sparing TRBC1+ cells in a mixed tumour model

Summary

- Targeting TCR beta constant chain isoforms may provide a generic approach to tackle the multiple subtypes of T cell-lymphoma
- AUTO4 clinical study in progress
- Development of a selective α TRBC2 binder for CAR T therapy
- Preclinical studies package demonstrating utility of α TRBC2 CAR *in vitro* and *in vivo*
 - Efficient CAR T proliferation upon target cell interaction *in vitro*
 - Significant IFN γ and IL2 secretion in response to target cell encounter *in vitro*
 - *In vivo* clearance of TRBC2 tumor in single and mixed tumor NSG models



Autolus


Solid Tumors

AUTO6NG – overcomes immune suppressive mechanisms in the tumor microenvironment (TME) and demonstrate preclinical anti-tumor activity in GD2-expressing solid tumors


AUTO6 designed to drive anti-tumor activity without neurotoxicity

AUTO6: GD2-targeted programmed T cell therapy in neuroblastoma


- Programmed CAR T clinical candidate
 - New binder design
 - Minimize on-target, off-tumor toxicity
 - Humanized to reduce immunogenicity
 - RQR8 safety switch
- Phase I trial in r/r Neuroblastoma conducted by CRUK in collaboration with UCL
- Autolus has the rights to clinical data and patents



Day 0



Day 28




MIBG: iodine-123-meta-iodobenzylguanidine

Targeting GD2 in indications outside Neuroblastoma


Screening for GD2 expression in multiple solid tumors

- GD2 is highly expressed in Osteosarcoma, Small Cell Lung Cancer (SCLC), Melanoma, and absent in Breast Cancer (TMA IHC)
- Detailed expression in SCLC was analyzed further in a larger set (25 SCLC patients' biopsies)
 - 68% GD2 expressing (homogenous and heterogenous)
 - 32% No GD2




Immune suppressive pathways in SCLC

Action in T Cell Synapse



Nikolaos Patsoukis et al, Front. Immunol., 11 April 2017




Action on Immune Cell Migration



Mariacarmela Santarpia et al, TCCR., Vol 4, No 6 (December 2015)

TGF- β status in a panel of SCLC biopsies

A strong pSMAD3 signal in these patients' biopsies




- Multiplex & multi-dimensional analysis by tissue cytometry
 - 28+ Markers Panel, analysed 10 SCLC biopsies
 - In depth characterization of the tumor cells and the components of the tumor microenvironment


AUTO6 is active *in vitro* – but not *in vivo* in a SCLC model

AUTO6 not efficacious *in vivo* in the H446 lung cancer model

AUTO6 active *in vitro*



AUTO6 not active *in vivo* in the same model




H446, a SCLC cell line expressing GD2
Average tumor size at infusion 20 mm³
Dose Administered: 7x10⁶ cells (IV)

AUTO6 CAR Ts actively traffic to tumor site

Trafficking is not culprit for lack of activity in the SCLC in vivo model

BLI – Day 10 after iv T cells



- AUTO6 cells co-localize with established subcutaneous H446 tumor
- However, CAR T cells do not appear to impact tumor cells
- Indicative of an inhibiting mechanism for the CAR Ts at the tumor site

Modular enhancements to drive AUTO6 efficacy

AUTO6NG designed to overcome tumor defenses, enhanced persistence and control

AUTO6

GD2 CAR

CAR design for optimal efficacy with minimal neurotoxicity

Safety Switch

To eliminate CAR Ts in the event of unexpected tox

dSHP2

To endow with checkpoint pathways resistance

Cytokine Signal

IL7R chimeric protein to improve persistence

dnTGFbRII* Receptor

To shield CAR Ts from immuno-suppressive signals


in vitro functional activity of AUTO6NG

Efficacy readily demonstrated *in vitro* in the H446 SCLC model

AUTO6NG without IL7R CCR*




AUTO6NG



- NT Control
- ▼ AUTO6-dSHP2-dTGFβR
- ◆ AUTO6-dSHP2-dTGFβR-CCR7

in vivo functional activity of AUTO6NG in a SCLC


Addition cell programming modules drive efficacy in the H446 SCLC model



Average tumor size pre-infusion 30 mm³
Dose Administered: 6x10⁶ CAR Ts IV




In vivo biomarkers analysis: IL7R CCR enhances CAR T expansion

CAR Ts readily detected by flow at time of euthanasia (day 34 post infusion)



- CAR Ts persisted and detected by flow cytometry at tumor site and the spleen (Day 34)
- A higher number of CAR Ts persisted with the IL7R CCR* module

In vivo Biomarkers analysis: IL7R CCR does not drive cytokine secretion



- NT Control
- AUTO6-dSHP2-dTGF β R
- ▲ AUTO6-dSHP2-dTGF β R-CCR7

Conclusion

AUTO6NG is highly active *in vitro* and *in vivo* in a SCLC tumor model

- AUTO6, a GD2 targeting CAR is clinical active in Neuroblastoma
- Our experimental data indicates that GD2 is an attractive SCLC CAR T target
- AUTO6 alone is not sufficient to drive *in vivo* efficacy in a SCLC mouse model and additional cell programming modules rendering the CAR T cells insensitive to TGF- β signalling and checkpoint inhibition (AUTO6NG) are required to drive efficacy
- The modules enabled AUTO6NG cells to persist and be detected by flow on day 34 post infusion, and this expansion enhancement was without increasing cytokines levels *in vivo*
- AUTO6NG will be clinically explored in GD2 expressing tumors



Solid Tumors


AUTO7 – Anti-PSMA humanized CAR T cell with improved persistence and resistance to tumor microenvironment for metastatic castration resistant prostate cancer (mCRPC)

Metastatic Castration-Resistant Prostate Cancer (mCRPC)

mCRPC is a large commercial market but has few effective treatment options

- Prostate cancer is the second most common male cancer in the world¹
 - affects ~1.3 million and kills >360,000 people each year
 - addressable patient population for 2nd line mCRPC is projected at c.80,000 patients*
- High unmet medical need²
 - patients diagnosed with localized disease usually treated with surgery or radiotherapy; however, many develop more advanced recurrent disease
 - patients with advanced prostate cancer are primarily treated with hormone ablation therapy, yet tumors ultimately progress to become castration resistant^{3,4,5}
 - once metastasized the 5-year survival rate falls to less than 30%
 - existing therapies are not curative, while cell based therapies have the potential to be curative

Estimated number of new cases worldwide



Biological hurdles to address in mCRPC therapy

Therapeutic approach needs to address a complex tumor microenvironment (TME)

- Opportunity for immunotherapy exists but multiple parameters need to be addressed due to the complex TME influencing the response to immunotherapy:
 - Late stage prostate cancers appear to be immunologically cold tumors with minimal T cell infiltrates¹
 - mCRPC have limited response to single-agent checkpoint inhibition; despite promising results in early trials, anti-CLTA4 Ipilimumab showed a marginal response in mCRPC patients^{2,3}
 - PD-1/PDL1 targeting antibodies have been equally inadequate^{4,5}
 - Tackling overproduction of TGF β could overcome the immunosuppressive effects of this cytokine in the TME⁶
 - Acidification of the TME play an important role in the efficacy of therapeutics that target tumors⁷
- An effective immunological approach will have to combine multiple modalities. Systemic combination using protein or small molecule drugs risks inducing severe and cumulative adverse events



AUTO7 strategy is a CAR T approach with modular cellular programming

Our goal is to generate resilient CAR T cells withstanding hostile tumor environment

Feature	Biology	Protein Module	Systemic Impact
Specific Targeting	Novel anti-PSMA binder	PSMA-CAR	No
Resilience to Checkpoint Blockade	Block inhibitory receptor signals	dSHP2	No
Avoid TME-driven inhibition	Block TGF β signal	dnTGF β RII	No
Support CAR T cell Survival	Promotes Jak/Stat signal	IL7R CCR	No
Induce Specific Immune Response	Activate immune response at the tumor site	Low level secreted IL-12	No – limited to immediate tumor environment

AUTO7 modules designed to tackle complex solid tumour environment

Technology toolbox utilized to overcome hostile TME



Modules delivered using gamma-retroviral vector

An ultra sensitive AUTO7 CAR directed to the PSMA ligand on mCRPC


Module	Function
α -PSMA CAR	Specific lysis of mCRPC
RapaCasp	Suicide gene
dnTGF β RII	Shield TGF β suppression
dSHP2	block checkpoint inhibition
IL7_CCR	Promote CAR T cell persistence
SS-IL12	Host Immune Cell Recruitment

- PSMA is one of the most commonly expressed genes found in prostate cancers
- PSMA pattern of expression can vary with the degree of differentiation, and heterogeneity of expression can vary within the same tumor¹


An ultra sensitive AUTO7 CAR directed to the PSMA ligand on mCRPC

In house binder campaign to develop a resilient anti-PSMA binder that is stable at low pH

Staining on frozen prostate tissues




Effect of pH on stability




An ultra sensitive AUTO7 CAR directed to the PSMA ligand on mCRPC

Humanized α -PSMA CAR in AUTO7 is effective at killing high and low density PSMA positive cells

PSMA CAR



Target cell line	PSMA expression levels by ABC
SupT1-NT	0
SupT1-PSMA _{low}	578
SupT1-PSMA _{high}	16500



dnTGFβRII module desensitizes AUTO7 T cells to TGFβ



Module	Function
α-PSMA CAR	Specific lysis of mCRPC
RapaCasp	Suicide gene
dnTGFβRII	Shield TGFβ suppression
dSHP2	block checkpoint inhibition
IL7_CCR	Promote CAR T cell persistence
SS-IL12	Host Immune Cell Recruitment

- TGFβ is an anti-inflammatory cytokine and is expressed at high levels in the TME¹
- T cells are key targets for TGFβ mediated immune suppression; tumor-derived TGFβ compromises T-cell mediated antitumor immunity²
- TGFβ signalling promotes the epithelial-to-mesenchymal transition associated with high-grade mCRPC malignancy³ and acts as an important mediator of metastasis to specific organ sites

dnTGFβRII module desensitizes AUTO7 T cells to TGFβ

- dnTGFβRII maintains AUTO7 T cells cytolytic function in presence of an inhibitory cytokine TGFβ

dnTGFβRII



dSHP2 module shields AUTO7 T cells from checkpoint inhibition


Module	Function
α -PSMA CAR	Specific lysis of mCRPC
RapaCasp	Suicide gene
dnTGF β RII	Shield TGF β suppression
dSHP2	block checkpoint inhibition
IL7_CCR	Promote CAR T cell persistence
SS-IL12	Host Immune Cell Recruitment

- SHP2 mediates signaling from many inhibitory receptors such as PD-1¹
- mCRPC are known to overexpress the PD-1 ligand, PD-L1²
- Antibody blockade of the PD-1/PD-L1 pathway in mCRPC has resulted in partial clinical responses³


dSHP2 module shields AUTO7 T cells from checkpoint inhibition

dSHP2 module maintains AUTO7 cytolytic function in the presence of PD1/PDL1 signalling

dSHP2



Checkpoint Resistance



IL7R CCR module induces AUTO7 T cell persistence


Module	Function
α -PSMA CAR	Specific lysis of mCRPC
RapaCasp	Suicide gene
dnTGF β RII	Shield TGF β suppression
dSHP2	block checkpoint inhibition
IL7R CCR	Promote CAR T cell persistence
SS-IL12	Host Immune Cell Recruitment

- CAR T cell persistence correlates with clinical efficacy^{1,2}
- CAR T programs to solid tumors report poor persistence; T cells require cytokine signaling to proliferate
- IL7R CCR delivers a constitutive IL7 signal to the CAR T cells and prolongs CAR T cell persistence


IL7R CCR module induces AUTO7 T cell persistence

IL7R CCR extends AUTO7 activity through multiple rounds of re-stimulation


Cytokine signal



CCR driven proliferation



Enhanced serial killing



- E:T 1:2, CART-cells weekly rechallenged with fresh SupT1-PSMA target cells

SS-IL12 module maintains potent adjuvant activity with no toxicity


Module	Function
α -PSMA CAR	Specific lysis of mCRPC
RapaCasp	Suicide gene
dnTGF β RII	Shield TGF β suppression
dSHP2	block checkpoint inhibition
IL7_CCR	Promote CAR T cell persistence
SS-IL12	Host Immune Cell Recruitment

- IL-12 is a potent adjuvant that promotes a powerful anti-tumor response when combined with checkpoint inhibitors and adoptive T cell transfer¹. However, IL-12 has been associated with systemic toxicity²
- Engineered version of the IL-12 (SS-IL12) has been designed to constrain expression while maintaining its adjuvant activity


SS-IL12 module maintains potent adjuvant activity with no toxicity

SS-IL12 module prevents toxicity by drastically reducing IL-12 secretion

Immune Adjuvant




IL-12 secretion




SS-IL12 module maintains potent adjuvant activity with no toxicity

SS-IL12 module prevents toxicity and maintains potent anti-tumor activity in a murine model

- Mouse B16.F10 cells in C57BL/6 mice




Mouse weight




- CAR
- ▲ CAR-2A-IL12
- ▼ CAR-SS-IL12


CAR



CAR-2A-IL12




CAR-SS-IL12



AUTO7 demonstrates potent activity against PCa in a xenograft model






Module	Function
α -PSMA CAR	Specific lysis of mCRPC
RapaCasp	Suicide gene
dnTGF β RII	Shield TGF β suppression
dSHP2	block checkpoint inhibition
IL7R CCR	Promote CAR T cell persistence
SS-IL12	Host Immune Cell Recruitment



AUTO7 demonstrates potent activity against PCa in a xenograft model

AUTO7 CAR T-cells completely eradicated PSMA positive tumor with no signs of toxicity

- Human PC3-PSMA prostate cancer model in NSG mouse



AUTO7 overcomes immunotherapeutic challenges for mCRPC

Modular CAR T with interchangeable components designed to overcome challenges in mCRPC

PSMA CAR

AUTO7 CAR T cells are highly potent in cytotoxicity assays against high and low PSMA expressing tumor cells

dSHP2

AUTO7 CAR T cells are desensitized to PD-1/PD-L1 mediated immunosuppression

dnTGF β RII

Truncated TGF β RII receptor shields AUTO7 CAR T cells from TGF β inhibitory effect

Cytokine signal

IL7R CCR module confers AUTO7 CAR T cells cytokine independent proliferation and improved ability to serially kill targets

Immune Adjuvant

SS-IL12 module in AUTO7 CAR T cells maintains potent adjuvant activity with no toxicity

Summary and Next Steps

Dr. Christian Itin

Chairman and CEO

Multiple clinical data points expected through 2020

Product	Indication	Target	Event
B Cell Malignancies			
AUTO1	Adult ALL	CD19	<ul style="list-style-type: none"> • Ph1 long-term follow up Q2 & Q4 2020 • Ongoing recruitment and dose last patient H1 2021
AUTO1NG	Pediatric ALL	CD19 & 22	<ul style="list-style-type: none"> • Start Ph1 H2 2020
AUTO3	DLBCL	CD19 & 22	<ul style="list-style-type: none"> • Decision on Ph2 Q3 2020 • Full Ph1 data H2 2020
AUTO3NG	DLBCL	CD19 & 22	<ul style="list-style-type: none"> • Ready to start Ph1 H2 2020, life cycle mgmt
Multiple Myeloma			
AUTO8	Multiple Myeloma	BCMA & CAR X	<ul style="list-style-type: none"> • Start Ph1 study H2 2020
T Cell Lymphoma			
AUTO4	TRBC1+ Peripheral TCL	TRBC1	<ul style="list-style-type: none"> • Ph1 interim data H1 2021
GD2+ Tumors			
AUTO6NG	Neuroblastoma; Melanoma; Osteosarcoma; SCLC	GD2	<ul style="list-style-type: none"> • Start Ph1 H1 2021
Allogeneic Approach			
Undisclosed	Undisclosed	Undisclosed	<ul style="list-style-type: none"> • Start Ph1 Q4 2020

Q&A


Dr. Christian Itin (Chairman and CEO)

Dr. Martin Pule (Founder & SVP, CSO)

Dr. Vijay Reddy (SVP, CMO)

Dr. Muhammad Al-Hajj (SVP, Head of Translational Research)

Dr. Mathieu Ferrari and Dr Marco Della Peruta



Thank you